Brain Research, 530 (1990) 358–360
Elsevier

Author Index

Addis, A., see Forloni, G.-L., 156
Altschuler, R.A., see López, I., 170
Amoroso, D., see Forloni, G.-L., 156
Angeretti, N., see Forloni, G.-L., 156
Aschner, M., Eberle, N.B., Miller, K. and Kimelberg, H.K., Interactions of methylmercury with rat primary astrocyte cultures: inhibition of rubidium and glutamate uptake and induction of swelling, 245

Barbosa, S.P., Saad, W.A., De Arruda Camargo, L.A., Renzi, A., De Luca, Jr., L.A., Fracasso, J.F. and Menani, J.V., Lesion of the anteroventral third ventricle region abolishes the beneficial effects of hypertonic saline on hemorrhagic shock in rats, 342
Barraco, R.A., El-Ridi, M.R. and Parizon, M., The adenosine analog, 5′-*N*-ethylcarboxamidoadenosine, exerts mixed agonist actions on cardiorespiratory parameters in the intact but not decerebrate rat following microinjections into the nucleus tractus solitarius, 54
Barth, D.S. and Di, S., The electrophysiological basis of epileptiform magnetic fields in neocortex, 35
Bartrup, J.T. and Stone, T.W., Activation of NMDA receptor-coupled channels suppresses the inhibitory action of adenosine on hippocampal slices, 330
Bartsch, U., see Blaugrund, E., 239
Baskys, A., Reynolds, J.N. and Carlen, P.L., NMDA depolarizations and long-term potentiation are reduced in the aged rat neocortex, 142
Berger, B., see Gaspar, P., 181
Biegon, A., see Gross-Isseroff, R., 312
Bisby, M.A. and Chen, S., Delayed Wallerian degeneration in sciatic nerves of C57BL/OIa mice is associated with impaired regeneration of sensory axons, 117
Black, S.E., see Kertesz, A., 40
Blaha, C.D., see Pfaus, J.G., 345
Blaugrund, E., Bartsch, U., Martini, R., Schachner, M. and Schwartz, M., Immunological evidence that the neural adhesion molecule L1 is expressed in fish brain and optic nerve: possible association with optic nerve regeneration, 239
Boadle-Biber, M.C., see Singh, V.B., 49
Bosler, O., see Saidi, H., 151
Bour, H., see Gagne, C., 197
Braak, E., see Ulfig, N., 176
Brodie, M.S., see Osmanovic, S.S., 287
Brown, A.G., see Maxwell, D.J., 215
Bugiani, O., see Giaccone, G., 325
Busija, D.W., see Shibata, M., 267

Calkoen, F., see Holstege, J.C., 130
Canevari, L., see Dagani, F., 261
Carlen, P.L., see Baskys, A., 142
Casagrande, V.A., see Claps, A., 126
Chen, S., see Bisby, M.A., 117
Choi, D.W., see Giffard, R.G., 138
Christie, W.M., see Maxwell, D.J., 215
Claps, A. and Casagrande, V.A., The distribution and morphology of corticogeniculate axons in ferrets, 126
Condemi, G., see Sica, A.L., 349
Connick, J.H., see Mally, J., 353
Consolo, S., see Forloni, G.-L., 156
Coote, J.H., see Lewis, D.I., 229
Crews, D., see Morris, Y.A., 339

Dagani, F., Ferrari, R. and Canevari, L., A pharmacological model for studying the role Na^+ gradients in the modulation of synaptosomal free $[Ca^{2+}]_i$ levels and energy metabolism, 261
Damsma, G., see Pfaus, J.G., 345
Dashwood, M.R., see Marks, S.A., 321
De Arruda Camargo, L.A., see Barbosa, S.P., 342
Del Cerro, S., Larson, J., Oliver, M.W. and Lynch, G., Development of hippocampal long-term potentiation is reduced by recently introduced calpain inhibitors, 91
De Li, C., see Lin, Q., 317
De Luca, Jr., L.A., see Barbosa, S.P., 342
De Mello, F.G., see Ventura, A.L.M., 301
Di, S., see Barth, D.S., 35
Di Lorenzo, P.M., Corticofugal influence on taste responses in the parabrachial pons of the rat, 73
Di Prisco, G.V., Wallén, P. and Grillner, S., Synaptic effects of intraspinal stretch receptor neurons mediating movement-related feedback during locomotion, 161
Dillon, K.A., see Gross-Isseroff, R., 312
Dragunow, M., Williams, M. and Faull, R.L.M., Haloperidol induces Fos and related molecules in intrastriatal grafts derived from fetal striatal primordia, 309
Dreifuss, J.J., see Raggenbass, M., 85
Dubois-Dauphin, M., see Raggenbass, M., 85

Eberle, N.B., see Aschner, M., 245
Elepfandt, A., see Traub, B., 105
El-Ridi, M.R., see Barraco, R.A., 54

Farlow, M.R., see Giaccone, G., 325
Faull, R.L.M., see Dragunow, M., 309
Febvret, A., see Gaspar, P., 181
Ferrari, R., see Dagani, F., 261
Fibiger, H.C., see Pfaus, J.G., 345
Forloni, G.-L., Angeretti, N., Amoroso, D., Addis, A. and Consolo, S., Decrease in [^{3}H]hemicholinium binding to high-affinity choline uptake sites in deafferented striatum: restoration by oxiracetam, 156
Fracasso, J.F., see Barbosa, S.P., 342
Frangione, B., see Giaccone, G., 325
Freed, W.J., see Simonds, G.R., 12
Furukawa, K., Yamana, K. and Kogure, K., Postischemic alterations of spontaneous activities in rat hippocampal CA1 neurons, 257

Gagne, C., Moyse, E., Kocher, L., Bour, H. and Pujol, J.F., Light-microscopic localization of somatostatin binding sites in the locus coeruleus of the rat, 196
Gandhi, M.R., see Sica, A.L., 349
Gaspar, P., Berger, B. and Febvret, A., Neurotensin innervation of the human cerebral cortex: lack of colocalization with catecholamines, 181
Gean, P.-W., The epileptiform activity induced by 4-aminopyridine in rat amygdala slices: antagonism by non-*N*-methyl-D-aspartate receptor antagonists, 251
Ghetti, B., see Giaccone, G., 325
Giaccone, G., Tagliavini, F., Verga, L., Frangione, B., Farlow, M.R., Bugiani, O. and Ghetti, B., Neurofibrillary tangles of the Indiana kindred of Gerstmann-Sträussler-Scheinker disease share antigenic determinants with those of Alzheimer disease, 325
Giffard, R.G., Monyer, H. and Choi, D.W., Selective vulnerability of cultured cortical glia to injury by extracellular acidosis, 138
Gilbey, M.P., see Marks, S.A., 321
Grillner, S., see Di Prisco, G.V., 161
Gross-Isseroff, R., Dillon, K.A., Israeli, M. and Biegon, A., Regionally selective increases in μ-opioid receptor density in the brains of suicide victims, 312

Harigaya, Y., see Shoji, M., 113
He, X., Schepelmann, K., Schaible, H.-G. and Schmidt, R.F., Capsaicin inhibits responses of fine afferents from the knee joint of the cat to mechanical and chemical stimuli, 147
Hirai, S., see Shoji, M., 113
Holstege, J.C. and Calkoen, F., The distribution of GABA in lumbar motoneuronal cell groups. A. quantitative ultrastructural study in rat, 130
Honoré, T., see Imperato, A., 223
Howell, J., see Kertesz, A., 40

Imperato, A., Honoré, T. and Jensen, L.H., Dopamine release in the nucleus caudatus and in the nucleus accumbens is under glutamatergic control through non-NMDA receptors: a study in freely-moving rats, 223
Israeli, M., see Gross-Isseroff, R., 312

Jensen, L.H., see Imperato, A., 223
Jones, S.L. and Light, A.R., Electrical stimulation in the medullary nucleus raphe magnus inhibits noxious heat-evoked *fos* protein-like immunoreactivity in the rat lumbar spinal cord, 335
Juiz, J.M., see López, I., 170

Kanje, M., see Sjöberg, J., 167

Karler, R., see Partlow, L.M., 102
Katz, M.J., see Price, R.L., 205
Kawarabayashi, T., see Shoji, M., 113
Kertesz, A., Polk, M., Black, S.E. and Howell, J., Sex, handedness, and the morphometry of cerebral asymmetries on magnetic resonance imaging, 40
Kimelberg, H.K., see Aschner, M., 245
Kocher, L., see Gagne, C., 196
Kogure, K., see Furukawa, K., 257

Larson, J., see Del Cerro, S., 91
Lasek, R.J., see Price, R.L., 205
Leffler, C.W., see Shibata, M., 267
Levine, J.D., see Sutters, K.A., 290
Lewis, D.I. and Coote, J.H., Excitation and inhibition of rat sympathetic preganglionic neurones by catecholamines, 229
Li, P., see Lin, Q., 317
Light, A.R., see Jones, S.L., 335
Lin, Q., De Li, C. and Li, P., Spinal cord substance P mediates carbachol-induced cardiovascular responses from the rostral ventrolateral medulla, 317
López, I., Juiz, J.M., Altschuler, R.A. and Meza, G., Distribution of GABA-like immunoreactivity in guinea pig vestibular cristae ampullaris, 170
Lund, R.D., see Yee, K.T., 121
Lynch, G., see Del Cerro, S., 91

Mally, J., Connick, J.H. and Stone, T.W., Chronic benzodiazepine treatment and cortical responses to adenosine and GABA, 353
Mangurian, L.P., see Walsh, R.J., 1
Marks, S.A., Stein, R.D., Dashwood, M.R. and Gilbey, M.P., [^{3}H]Prazosin binding in the intermediolateral cell column and the effects of iontophoresed methoxamine on sympathetic preganglionic neuronal activity in the anaesthetized cat and rat, 321
Martini, R., see Blaugrund, E., 239
Maxwell, D.J., Christie, W.M., Short, A.D. and Brown, A.G., Direct observations of synapses between GABA-immunoreactive boutons and muscle afferent terminals in lamina VI of the cat's spinal cord, 215
Menani, J.V., see Barbosa, S.P., 342
Meza, G., see López, I., 170
Miaskowski, C., see Sutters, K.A., 290
Miceli, D., see Rio, J.P., 96
Miller, J.F. and Proudfit, H.K., Antagonism of stimulation-produced antinociception from ventrolateral pontine sites by intrathecal administration of α-adrenergic antagonists and naloxone, 20
Miller, K., see Aschner, M., 245
Monyer, H., see Giffard, R.G., 138
Morris, Y.A. and Crews, D., The effects of exogenous neuropeptide Y on feeding and sexual behavior in the red-sided garter snake (*Thamnophis sirtalis parietalis*), 339
Moyse, E., see Gagne, C., 196

Nakagawa, S., see Nakamura, H., 108
Nakamura, H., Saheki, T. and Nakagawa, S., Differential cellular localization of enzymes of L-arginine metabolism in the rat brain, 108
Nomikos, G.G., see Pfaus, J.G., 345

Ohm, T.G., see Ulfig, N., 176
Oliver, M.W., see Del Cerro, S., 91
Onaivi, E.S., see Singh, V.B., 49
Osmanovic, S.S., Shefner, S.A. and Brodie, M.S., Functional significance of the apamin-sensitive conductance in rat locus coeruleus neurons, 287

Parizon, M., see Barraco, R.A., 54
Partlow, L.M., Karler, R. and Turkanis, S.A., Whole-cell voltage-clamp study of sodium current in neuroblastoma cells: effects of inhibition of neurite outgrowth by colchicine, 102
Peyrichoux, J., see Rio, J.P., 96
Pfaus, J.G., Damsma, G., Nomikos, G.G., Wenkstern, D.G., Blaha, C.D., Phillips, A.G and Fibiger, H.C., Sexual behavior enhances central dopamine transmission in the male rat, 345
Phan, T.-H., see Singh, V.B., 49
Phillips, A.G., see Pfaus, J.G., 345
Polk, M., see Kertesz, A., 40
Pool, C.W., see Ulfig, N., 176
Posner, B.I., see Walsh, R.J., 1
Price, R.L., Lasek, R.J. and Katz, M.J., Internal axonal cytoarchitecture is shaped locally by external compressive forces, 205
Proudfit, H.K., see Miller, J.F., 20
Pujol, J.F., see Gagne, C., 196

Raggenbass, M., Vozzi, C., Tribollet, E., Dubois-Dauphin, M. and Dreifuss, J.J., Thyrotropin-releasing hormone causes direct excitation of dorsal vagal and solitary tract neurones in rat brainstem slices, 85
Rao, K., see Yee, K.T., 121
Rehfeld, J.F., Van Solinge, W.W., Tos, M. and Thomsen, J., Gastrin, cholecystokinin and their precursors in acoustic neuromas, 235
Renzi, A., see Barbosa, S.P., 342
Repérant, J., see Rio, J.P., 96
Reynolds, J.N., see Baskys, A., 142
Rio, J.P., Réperant, J., Miceli, D. and Peyrichoux, J., Reorganization of GABAergic synapses in the viper optic tectum following retinal deafferentation, 96

Saad, W.A., see Barbosa, S.P., 342
Saheki, T., see Nakamura, H., 108
Saidi, H. and Bosler, O., Serotonin reinnervation of the rat organum vasculosum laminae terminalis (OVLT) after 5,7-dihydroxytryptamine deafferentation, 151
Schachner, M., see Blaugrund, E., 239
Schaible, H.-G., see He, X., 147
Schepelman, K., see He, X., 147
Schmidt, R.F., see He, X., 147
Schwartz, M., see Blaugrund, E., 239
Sessler, F.M., see Stone, E.A., 295
Shefner, S.A., see Osmanovic, S.S., 287
Shibata, M., Leffler, C.W. and Busija, D.W., Cerebral hemodynamics during cortical spreading depression in rabbits, 267
Shoji, M., Hirai, S., Harigaya, Y., Kawarabayashi, T. and Yamaguchi, H., The amyloid β-protein precursor is localized in smooth muscle cells of leptomeningeal vessels, 113
Short, A.D., see Maxwell, D.J., 215
Sica, A.L., Siddiqi, Z.A., Gandhi, M.R. and Condemi, G., Evidence for central patterning of sympathetic discharge in kittens, 349
Siddiqi, Z.A., see Sica, A.L., 349
Simonds, G.R. and Freed, W.J., Effects of intraventricular substantia nigra allografts as a function of donor age, 12
Singh, V.B., Onaivi, E.S., Phan, T.-H. and Boadle-Biber, M.C., The increases in rat cortical and midbrain tryptophan hydroxylase activity in response to acute or repeated sound stress are blocked by bilateral lesions to the central nucleus of the amygdala, 49
Sjöberg, J. and Kanje, M., Effects of repetitive conditioning crush lesions on regeneration of the rat sciatic nerve, 167
Slater, N.T., see Smith, R.D., 275
Smetanka, A.M., see Yee, K.T., 121
Smith, R.D., Turek, F.W. and Slater, N.T., Bicuculline and picrotoxin block phase advances induced by GABA agonists in the circadian rhythm of locomotor activity in the golden hamster by a phaclofen-insensitive mechanism, 275
Stein, R.D., see Marks, S.A., 321
Stone, E.A., Sessler, F.M. and Weimin, L., Glial localization of adenylate-coupled β-adrenoceptors in rat forebrain slices, 295
Stone, T.W., see Bartrup, J.T., 330
Stone, T.W., see Mally, J., 353
Sutters, K.A., Miaskowski, C., Taiwo, Y.O. and Levine, J.D., Analgesic synergy and improved motor function produced by combinations of μ-δ- and μ-κ-opioids, 290

Tagliavini, F., see Giaccone, G., 325
Taiwo, Y.O., see Sutters, K.A., 290
Thomsen, J., see Rehfeld, J.F., 235
Tos, M., see Rehfeld, J.F., 235
Traub, B. and Elepfandt, A., Sensory neglect in a frog: evidence for early evolution of attentional processes in vertebrates, 105
Tribollet, E., see Raggenbass, M., 85
Turek, F.W., see Smith, R.D., 275
Turkanis, S.A., see Partlow, L.M., 102

Ulfig, N., Braak, E., Ohm, T.G. and Pool, C.W., Vasopressinergic neurons in the magnocellular nuclei of the human basal forebrain, 176

Van Solinge, W.W., see Rehfeld, J.F., 235
Ventura, A.L.M. and De Mello, F.G., D_1 dopamine receptors in neurite regions of embryonic and differentiated retina are highly coupled to adenylate cyclase in the embryonic but not in the mature tissue, 301
Verga, L., see Giaccone, G., 325
Vozzi, C., see Raggenbass, M., 85

Wallén, P., see Di Prisco, G.V., 161
Walsh, R.J., Mangurian, L.P. and Posner,

B.I., The distribution of lactogen receptors in the mammalian hypothalaus: an in vitro autoradiographic analysis of the rabbit and rat, 1
Weimin, L., see Stone, E.A., 295
Wenkstern, D.G., see Pfaus, J.G., 345
Williams, M., see Dragunow, M., 309

Yamaguchi, H., see Shoji, M., 113
Yamana, K., see Furukawa, K., 257
Yee, K.T., Smetanka, A.M., Lund, R.D. and Rao, K., Differential expression of class I and class II major histocompatibility complex antigen in early postnatal rats, 121

(contents continued)

G. Giaccone, F. Tagliavini, L. Verga, B. Frangione, M.R. Farlow, O. Bugiani and B. Ghetti (Italy, U.S.A.) 325
Activation of NMDA receptor-coupled channels suppresses the inhibitory action of adenosine on hippocampal slices
J.T. Bartrup and T.W. Stone (U.K.) 330
Electrical stimulation in the medullary nucleus raphe magnus inhibits noxious heat-evoked *fos* protein-like immunoreactivity in the rat lumbar spinal cord
S.L. Jones and A.R. Light (U.S.A.) 335
The effects of exogenous neuropeptide Y on feeding and sexual behavior in the red-sided garter snake (*Thamnophis sirtalis parietalis*)
Y.A. Morris and D. Crews (U.S.A.) 339
Lesion of the anteroventral third ventricle region abolishes the beneficial effects of hypertonic saline on hemorrhagic shock in rats
S.P. Barbosa, W.A. Saad, L.A. de Arruda Camargo, A. Renzi, L.A. De Luca, Jr., J.F. Fracasso and J.V. Menani (Brazil) 342
Sexual behavior enhances central dopamine transmission in the male rat
J.G. Pfaus, G. Damsma, G.G. Nomikos, D.G. Wenkstern, C.D. Blaha, A.G. Phillips and H.C. Fibiger (Canada) 345
Evidence for central patterning of sympathetic discharge in kittens
A.L. Sica, Z.A. Siddiqi, M.R. Gandhi and G. Condemi (U.S.A.) 349
Chronic benzodiazepine treatment and cortical responses to adenosine and GABA
J. Mally, J.H. Connick and T.W. Stone (U.K.) 353

Author Index 358